

## Improved hodograph method to identify field-line resonances in ground magnetometer data

# Hideaki Kawano[1]; Viacheslav Pilipenko[2]; Satoko Takasaki[3]; Kiyohumi Yumoto[4]

[1] Earth and Planetary Sci., Kyushu Univ.; [2] IKI; [3] NIPR; [4] Space Environ. Res. Center, Kyushu Univ.

Where the frequency of a wave coming from outside the magnetosphere matches the eigen-frequency of a magnetospheric field line, the field line resonance (FLR) causes the field line in question to eigen-oscillate. Thus generated eigen-oscillations are useful for us to estimate the magnetospheric plasma mass density from the ground, because the eigen-frequency (FLR frequency below) decreases with increasing plasma mass along the field line.

However, FLR frequencies are often difficult to identify in the ground magnetometer data, because different kinds of waves with large amplitudes are often superposed onto the FLR signal and mask the FLR signal.

As countermeasures to this problem, methods called amplitude-ratio method and cross-phase method have been used; these methods take the difference between the data from two magnetometers that are latitudinally separated by an order of 100km. However, a problem here is that the two methods can yield different values of the FLR frequency from the same dataset. Another problem is that neither method can provide an estimation error (error bar) of the estimated FLR frequency.

Hodograph method is a solution to the former problem, because it merges the amplitude-ratio method and the cross-phase method into one method. Furthermore, it can estimate the resonance width. However, the resonance width is assumed to be a constant of latitude in the hodograph method, while in nature the resonance width can be a function of latitude. Furthermore, the hodograph method is still unable to provide an estimation error of the FLR frequency.

In this paper we intend to improve the hodograph method to solve the above-stated problems. That is, we introduce a latitude dependence of the resonance width into the hodograph method, and we also introduce into the method a framework to estimate errors. We will first test the validity of these modifications by applying the modified method to simulated datasets.