

地上-衛星観測による磁気嵐時の中緯度領域における電離圏電場について

辻 裕司 [1]; 西村 幸敏 [2]; 新堀 淳樹 [3]; 菊池 崇 [4]; 堀 智昭 [5]; 松岡 彩子 [6]; 長妻 努 [7]

[1] 名大・理・素粒子宇宙; [2] 名大 STEL; [3] 名大・太陽地球環境研究所; [4] STE 研; [5] STE 研; [6] 宇宙研; [7] NICT

Storm-time electric fields in the mid-latitude ionosphere derived from the ground and satellite observations

Yuji Tsuji[1]; Yukitoshi Nishimura[2]; Atsuki Shinbori[3]; Takashi Kikuchi[4]; Tomoaki Hori[5]; Ayako Matsuoka[6]; Tsutomu Nagatsuma[7]

[1] Particle and Astrophysical Sci., Nagoya Univ.; [2] STEL, Nagoya University; [3] Solar-Terrestrial Environment Laboratory, Nagoya Univ.; [4] STEL; [5] Solar-Terrestrial Environment Laboratory, Nagoya Univ.; [6] ISAS/JAXA; [7] NICT

In order to clarify a quantitative relationship between ground magnetic field variations and ionospheric electric fields in middle latitudes during a storm, we deduced the ionospheric electric fields from magnetic field variations measured on the ground and compared the deduced electric fields with in-situ electric fields observed by the Akebono satellite mapped onto the ionosphere. During the storm on February 21-22, 1994, with a minimum SYM-H value of -152 nT, dawn-dusk electric fields (4-6 mV/m) intensified in the dusk sector of the inner magnetosphere (L is less than 4). Near the end of the storm main phase (0100-0140 UT on February 22), the satellite passed over the North America in the dusk sector (\sim 1600-1800 magnetic local time) from 73 to 49 deg in invariant latitude (ILAT) where many magnetometers were installed. During this period, the satellite observed that the poleward electric field varied from negative to positive values at ILAT \sim 65.5 deg. The electric field signature indicates that the center of region-1 field-aligned currents was located at this latitude. Subsequently, the electric field showed two peaks at ILAT \sim 62-63 and 55 deg. The former and the later peaks correspond to large poleward electric fields for the auroral oval and SAPS/SAID, respectively. On the other hand, the electric fields deduced from ground magnetic field variations were consistent with those obtained from the satellite except for the auroral oval. In particular, it is noted that we could derive the enhanced electric fields in middle latitudes which corresponded to SAPS/SAID. These results suggest that it is possible to obtain the global distribution of ionospheric electric fields in middle and subauroral latitudes related to overshielding and SAPS/SAID from worldwide magnetometer networks.