

全球GNSS-TEC観測データベースを用いた磁気嵐時の電離圏擾乱に関する研究

新堀 淳樹 [1]; 大塚 雄一 [2]; 惣宇利 卓弥 [3]; 津川 卓也 [4]; 西岡 未知 [4]
[1] 名大・宇地研; [2] 名大宇地研; [3] 名大 ISEE; [4] 情報通信研究機構

Study of storm-time ionospheric disturbances using global GNSS-TEC observation database

Atsuki Shinbori[1]; Yuichi Otsuka[2]; Takuya Sori[3]; Takuya Tsugawa[4]; Michi Nishioka[4]
[1] ISEE, Nagoya Univ.; [2] ISEE, Nagoya Univ.; [3] ISEE, Nagoya Univ.; [4] NICT

The electron density distribution in the ionosphere is determined by the chemical and dynamical processes in addition to the solar extreme ultraviolet (EUV) ionization. Since the ionospheric electrons interact with electric fields of electromagnetic waves, the ionosphere gives a significant influence on the ionospheric transmissions with satellites for communications, positioning, and navigation. The ionosphere changes greatly in association with the activities of the sun, magnetosphere, and the lower atmosphere, and their ionospheric variations interfere with satellite communication and enhance satellite positioning errors in the Global Navigation Satellite System (GNSS), including GPS. The most major factor of the positioning errors is the propagation delay of GNSS signals in the ionosphere. Therefore, a global total electron content (TEC) map or ionospheric model is required to investigate global ionospheric variations associated with geomagnetic storms and lower atmospheric activities and to evaluate the ionospheric delay for GNSS positioning. A scale of the ionospheric variations (traveling ionospheric disturbances (TIDs), equatorial plasma bubbles (EPB), auroral ionospheric irregularity) is in a range between several 100 km and 1000 km. A storm enhanced density (SED) phenomenon observed in the mid- and low-latitude ionosphere during the main phase of geomagnetic storms appears with a wide longitudinal extent. In order to clarify generation and propagation mechanisms of these ionospheric phenomena and to investigate their effects on the GNSS positioning, dense and wide-coverage ionospheric observations and the corresponding TEC data are needed. Under these backgrounds, we have built a global TEC database (absolute TEC, detrended TEC, and ROTI) with high time and spatial resolutions and developed a TEC data analysis system in a corroboration with NICT and the IUGONET (Inter-university Upper atmosphere Global Observation NETWORK) and PWING (Study of dynamical variation of particles and waves in the inner magnetosphere using ground-based network observations) projects. The two-dimensional maps of absolute TEC, detrended TEC, and ROTI have been open on the TEC database websites (<http://stdb2.isee.nagoya-u.ac.jp/GPS/GPS-TEC/index.html> and <http://seg-web.nict.go.jp/GPS/DRAWING-TEC/>). For the development of the TEC database, we have collected all the available GNSS receiver data in the world. The number of the GNSS stations reaches more than 8500 in January 2019. These GNSS data are provided by the International Geoscience Services (IGS), the University NAVSTAR Consortium (UNAVCO), Scripps Orbit and Permanent Array Center (SOPAC), and other global and regional data centers (more than 50 data providers in all). In this talk, we introduce an overview of recent new scientific results of temporal and spatial evolutions of storm-time ionospheric disturbances (longitudinal variation of the midlatitude trough [Shinbori et al., 2018], plasmaspheric erosion [Obana et al., 2019], longitudinal extent of the SED plume [Sori et al., 2019], midlatitude plasma bubbles [Sori et al., 2019]) using the global GNSS-TEC database.