

R005-22

Zoom meeting C : 11/1 PM2 (15:45-17:30)
16:30-16:45

Variations in the D-region ionosphere observed in fireballs using VLF/LF transmitter signals.

#Takeru Suzuki¹⁾, Hiroyo Ohya²⁾, Fuminori Tsuchiya³⁾, Kazuo Shiokawa⁴⁾, Hiroyuki Nakata²⁾

¹⁾Science and Engineering, Chiba University, ²⁾Engineering, Chiba Univ., ³⁾Planet. Plasma Atmos. Res. Cent., Tohoku Univ., ⁴⁾ISEE, Nagoya Univ.

Meteors and fireballs are known to ionize the D-region and lower E-region ionospheres at 80-120 km heights [Davies, 1966]. The fireballs are meteors that the magnitude of brightness is larger than -4 based on the IAU (International Astronomy Union) definition. TID (traveling ionospheric disturbance) associated with the Chelyabinsk meteoroid in Russia was reported based on GPS-TEC (total electron content) observations [Perevalova et al., 2015]. The amplitude of the TEC variations was 0.07-0.5 TECU, and the period was 10 minutes. The epicenter of the TID was airburst point at 20-30 km heights of the meteoroid, and the velocities were 250-660 m/s. However, few quantitative studies for the D-region ionosphere associated with meteors and fireballs have been reported. In this study, we investigate the variations in the D-region ionosphere during a fireball occurred in Hokkaido, using VLF (very low frequency, 3-30 kHz) / LF (low frequency, 30-300 kHz) transmitter signals. The VLF/LF transmitter signals are reflected in the D-region ionosphere and are sensitive for variations in electron density in the lower ionosphere. The transmitters used in this study were JY40kHz (Fukushima, Japan, 37.37° N, 140.85° E), JY60kHz (Saga, Japan, 33.47° N, 130.18° E), and JI (Miyazaki, Japan, 22.2 kHz, 32.05° N, 130.82° E). The receiver was located at RKB (Rikubetsu, Japan, 43.45° N, 143.77° E). Periodic variations of 100-200 s were identified by a wavelet transformation of the signal intensities for the JY40kHz-RKB, JY60kHz-RKB, and JI-RKB paths at about five minutes (12:01 UT) after the fireball. The phases of the JY40kHz-RKB and JY60kHz-RKB paths had similar periods of 100-200 s at that time. We consider that these variations of intensity and phase were caused by the D-region variations due to acoustic waves in the atmosphere excited by the fireball. If the acoustic waves were excited at the luminous point (118 km altitude), the VLF/LF reflection point (90 km altitude) or vanishing end point (25 km altitude) of the fireball, the propagation times of the acoustic waves from the excited point to the LF reflection point at 90 km height over RKB were calculated to be 138 s, 126 s, or 311 s, respectively. The arrival time (311 s) of the acoustic waves excited from the vanishing point at the 25 km altitude was in good agreements with the onset of the VLF/LF variations with the period of 100-200 s. The acoustic waves excited at the vanishing end point propagate upward and may cause the variations in electron density in the D-region ionosphere. We will also show results of another fireball event over Kanto region, Japan, at 17:32:02 UT on 1 July, 2020, and discuss the fireball effects on the D-region ionosphere.