

R005-09

B会場：9/24 PM2 (15:45-18:15)

16:15~16:30

#古舘 千力<sup>1)</sup>, 西山 尚典<sup>2)</sup>, 津田 卓雄<sup>1)</sup>, 小川 泰信<sup>2)</sup>, 堤 雅基<sup>2)</sup>, Partamies Noora<sup>3)</sup>, 野澤 悟徳<sup>4)</sup>, Sigernes Fred<sup>3)</sup>  
(<sup>1)</sup> 電通大, (<sup>2)</sup> 極地研, (<sup>3)</sup> University Centre in Svalbard, (<sup>4)</sup> 名大・宇地研

## Atmospheric wave variability in the upper mesosphere based on ground-based observations of OH airglows (~1.3 $\mu$ m) in Longyearbyen

#Senri Furutachi<sup>1)</sup>, Takanori Nishiyama<sup>2)</sup>, Takuo Tsuda<sup>1)</sup>, Yasunobu Ogawa<sup>2)</sup>, Masaki Tsutsumi<sup>2)</sup>, Noora Partamies<sup>3)</sup>, Satonori Nozawa<sup>4)</sup>, Fred Sigernes<sup>3)</sup>

(<sup>1)</sup>University of Electro-Communications, (<sup>2)</sup>National Institute of Polar Research, (<sup>3)</sup>University Centre in Svalbard, (<sup>4)</sup>Institute for Space-Earth Environment Research, Nagoya University

The energy and momentum of the atmospheric waves, such as atmospheric gravity waves, tides, and planetary waves, can drive the zonal and/or meridional winds in the mesosphere and lower thermosphere (MLT), which are considered to be closely related to the general circulation in the whole atmosphere. Thus, it is important to observe the activities of such atmospheric waves in the MLT region, for more precise modeling research. Many observations of such atmospheric waves have been done for many years. In particular, recent polar-orbit satellites have been providing global observations. On the other hand, in addition to such satellites, higher resolution observations by ground-based observations are also important to resolve smaller- or shorter-scale waves. For example, tides with higher frequencies (8- and 6-hours periods) in polar regions have not been fully studied and poorly understood

As for the temperature observations from the ground, the rotational temperature of OH airglow has been extensively investigated in the past over 60 years. Recently, short-wavelength infrared OH airglow observations using InGaAs FPA (Focal Plane Array) have been reported. The OH airglow intensity in this region is stronger than that in the visible subrange, and thus more advanced airglow observations (e.g., with higher time resolution) can be expected. However, observations using the short-wavelength infrared OH bands are still limited so far, especially in the polar region. In addition, auroral contamination is one of the difficulties in airglow observations in the polar region. For example, there is a report that temperature measurements using the OH (3,1) band (~1.5  $\mu$  m) at Syowa Station, Antarctic (69.0 °N, 39.6 °N) included an underestimation of the temperature possibly by ~40 K due to auroral contamination. Therefore, a more robust method is needed for OH temperature observations in the polar region.

In this research, we propose spectroscopic observations of the OH airglow in ~1.3  $\mu$  m, which would be expected to be more robust to such auroral contamination. We have developed a brand new Near-Infrared Aurora and airglow Spectrograph-2 (NIRAS-2). It is an imaging spectrograph with InGaAs 2D FPA, which has a wide FOV of 55 degrees with a resolution of 0.11 degrees, and its wavelength resolution is variable with combinations of three slits, 30-, 60-, and 90-  $\mu$  m, and two volume phase holographic gratings, 950- and 1500-lpmm. OH airglow observations are mainly performed using the low-dispersion 950-lpmm grating with 60-  $\mu$  m slits. The wavelength range is from 1195 to 1350 nm, targeting the OH (7,4) and (8,5) bands and O<sub>2</sub> IR band with a spectral resolution of 1.1 nm. The preliminary test observation in NIPR, Tokyo, showed that the signal-noise ratio is better than its predecessor, NIRAS. The temperature data, estimated from the OH (8,5) band (~1.3  $\mu$  m) in the 10-minute integration, were well comparable with those from NRLMSISE-00. The estimated temperature error was a few K, which is significantly better than that from NIRAS.

NIRAS-2) was installed at The Kjell Henriksen Observatory (KHO), Longyearbyen (78.1 °N, 16.0 °N) in late November 2022. Continuous 24-hour observations of the OH bands were made with 30-second exposures from November 23 to December 26, 2022, and then the OH observations were also routinely done for continuous two weeks in every month from January to March 2023. OH rotational temperatures derived from the 30-minute integrations are in a good agreement with the OH (6,2) rotational temperatures obtained from the spectrometer of the University Centre in Svalbard and the temperature obtained from Aura/Microwave Limb Sounder. Moreover, we are working on data analysis to investigate the activities of atmospheric waves, mainly atmospheric tides and gravity waves. In the preliminary analysis, we found clear 6- and 8-hour wave activities in both the OH (7,4) and OH (8,5) bands data with 5-min resolutions. These results would be similar to the previous results obtained at Eureka (80 °N) [Oznovich et al., 1997]. For further investigation, the results from the temperature observations will be then compared with wind observations from the co-located Nippon/Norway Svalbard Meteor Radar (NSMR) [Hall et al., 2002]. In the presentation, we will show these results and give a more detailed discussion.